



**Specifications For Proper Pesticide Use**  
**Read and Follow Label Directions**



<b>Product</b>	<b>REI (Re Entry Interval - hours)</b>	<b>PHI (Pre Harvest Interval - days)</b>	<b>Ground Application Water Volume (gallons)</b>	<b>Rainfast (hours)</b>
<b>Herbicides</b>				
Assure II, Targa	12	45	Min 10	1
Betamix, D-P Mix, Phen-Des	24	75	10 - 20	6
Betanex, Des, Alphanex	24	75	10 -20	6
Dual Magnum	24	60	Min 10	0
Eptam / Eptek	12	---	10 - 50	0
Fargo (Granular)	12	---	-----	0
Fargo 4EC (liquid)	12	---	Min 10	0
Glyphosate (RoundUp)	12	---	3 - 40	*** (see reference below)
Gramoxone Max / Paraquat	12	---	Min 10	4 - 6
Herbicide - 273	48	60 (DAE - Days After Emergence)	10 - 40	1/2 or 30 minutes
Nortron, Etho SC, Ethotron	12	---	10 - 60	6 (foliar) 0 (soil)
Outlook, Propel, Establish	12	60	5 or more	0
Poast	12	60	5 - 20	1
Progress, Des-Phen-Etho	48	75	10 -20	6
RoNeet	12	---	10 - 50	0
Select Max, Select, Arrow	24	40	Min 10 with broadleaf tank mix	1
Volunteer, Trigger, Clethodim, Section				
Stinger, Clopyr Ag, Spur, Garrison	12	45	10 or more	6
Treflan	12	---	5 - 40	0
UpBeet	4	60	Min 10	6

**WARNING:** This information is based on pesticides registered with EPA in January 2007. Sugarbeet treated with a non-registered pesticide or other chemical may have illegal residues and violate the grower 5 year agreement and will not be accepted by American Crystal Sugar Company for delivery. When applying pesticides after 2007 check labels registered with EPA for proper pesticide use recommendations.



## Specifications For Proper Pesticide Use



<b>Product</b>	<b>REI (Re Entry Interval - hours)</b>	<b>PHI (Pre Harvest Interval - days)</b>	<b>Ground Application Water Volume (gallons)</b>	<b>Rainfast (hours)</b>
<b>Insecticides</b>				
Asana	12	21	Min 2	avoid applying where surface water is present
Counter	48 - 72	110	-----	0
Diazinon	24	14	Use sufficient water for uniform coverage	avoid applying where surface water is present
Dibrom 8	48	2	1 - 5	+ + + (see reference below)
Lannate	48	21	Use sufficient water for uniform coverage	avoid applying where surface water is present
Lorsban 15G, Chlorpyrifos 15G, Nufos 15G	24	7	-----	0
Lorsban 4E, Chlorpyrifos 4E, Nufos 4E, Govern 4E, Warhawk, Whirlwind	24	30	10 - 30	0
Methyl parathion	96 - 120	20	Min 3	6
Mustang Max	12	50	Min 10 (foliar) Min 3-5 (soil applied)	avoid applying where surface water is present (foliar)
Sevin 4F	12	28	Use sufficient water for uniform coverage	avoid applying where surface water is present
Temik	48	90	-----	0
Thimet	48 - 72	30	-----	0
<b>Fungicides</b>				
Agri Tin	48	7	At least 15	avoid applying where surface water is present
Amistar, Quadris	4	0	sufficient water to ensure thorough coverage of foliage	avoid applying where surface water is present
Eminent	12	14	5 or more	avoid applying where surface water is present
Enable 2F	12	14	20	1
Gem	12	21	Min 10	avoid applying where surface water is present
Headline	12	7	sufficient water to ensure thorough coverage of foliage	avoid applying where surface water is present
Mancozeb	24	14	Min 2	avoid applying where surface water is present
Maneb	24	14	Min 10	avoid applying where surface water is present
Sulfurs	24	0	Min 5	avoid applying where surface water is present
Supertin (Liquid and 80 WP)	48	7	At least 15	avoid applying where surface water is present
Topsin	12	21	Use sufficient water for uniform coverage	avoid applying where surface water is present

\*\*\* Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control. (6 hours usually adequate)

+ + + Do not apply within 24 hours following rainfall or irrigation, or in areas where intense or sustained rainfall is forecasted to occur within 24 hours following application