



2020 Layby Herbicide Control Options

Post-emergence control options for glyphosate resistant Waterhemp are very limited and are becoming less effective. Therefore, a layby application approach with chloroacetamide herbicides such as Outlook, Warrant, or Dual Magnum is necessary to continue to provide a barrier against Waterhemp emergence. Which option you choose should be based upon timing, likelihood of rainfall and finally if you used a PRE or PPI soil applied herbicide at planting. All layby herbicides need to be applied after the first 2 true leaves have emerged and the stand has been established. As with any pest control strategy, timing of application is more important than what herbicide you choose to use. Please consult your Agriculturalist for further information and Always Follow Label Directions.

Layby Herbicide: Soil applied herbicide that is applied after crop stand has been established (after 2 true leaf stage in sugarbeet). All layby applications need rainfall to activate the herbicide.

Herbicide	Rate	Crop Injury Risk	Notes
Outlook	<p>Single - 18oz application</p> <p>Split application 12 oz fb 12oz</p>	<p>Higher for single app</p> <p>Low for Split</p>	<ul style="list-style-type: none"> • Good Waterhemp control • Good Lambsquarters control • No Kochia control • Cannot replant to sugarbeet • Injury greater on small beets and course soil types • Very little moisture needed for activation $\approx \frac{1}{4}$" • 60 day PHI 2-8 leaf
Dual Magnum or generic S-metolachlor	<p>Single – 1.3pt/A course texture soil or up to 1.6pt/A fine texture soil</p> <p>Split application 1pt/A fb 1pt/A</p>	<p>Higher for single app</p> <p>Low for Split</p>	<ul style="list-style-type: none"> • Good Waterhemp control • Fair Lambsquarters control • Poor Kochia control • Only use S-metolachlor formulations for best results • Beware: metolachlor products are cheap but not labeled in sugarbeet • Needs $\frac{1}{2}$" rainfall to activate • 60 day PHI
Warrant	<p>Single 1.5-2qts/A</p> <p>Split 2.5pt-3pt fb 2.5pt -3pt/A</p>	<p>Higher for single app</p> <p>Low for Split</p>	<ul style="list-style-type: none"> • Good Waterhemp control • Fair Lambsquarter control • Poor Kochia control • Cannot replant to sugarbeet • Longest residual of any of the layby options • Needs minimum of $\frac{3}{4}$" of rainfall to activate • 70 day PHI

Additional Notes for all Layby Herbicides

- Layby herbicides must be applied to soil with good coverage to work properly
- Layby herbicides should be applied with 15 gallons of water and medium to course nozzles if sprayed alone
- Use at least 10 gallons of water if tank mixing layby with only glyphosate and AMS
- Use at least 12 gallons of water if tank mixing layby with glyphosate, AMS, and another broadleaf herbicide like Stinger, Betamix, ethofumesate, Upbeet, or an HSMOC surfactant
- Layby application can be delayed if a 5pt or greater rate of ethofumesate was used as a PRE or PPI. However, ethofumesate alone is not a season long Waterhemp control strategy. Use split application rates of layby herbicides where you applied ethofumesate, preemergence.
- Application with temperatures over 85F are more likely to cause injury